

An exact self-consistent gravitational shock wave in semiclassical gravity

Carlos O. Lousto*

*Department of Physics, University of Utah
201 JBF, Salt Lake City, UT 84112, USA*

Francisco D. Mazzitelli†

*Departamento de Física and IAFE
Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires
Ciudad Universitaria, Pabellón I, (1428) Buenos Aires, Argentina
(February 7, 2008)*

Abstract

We find a self-consistent pp-gravitational shock wave solution to the semi-classical Einstein equations resulting from the $1/N$ approach to the effective action. We model the renormalized matter stress-energy-momentum tensor by N massless scalar fields in the Minkowski vacuum plus a classical particle. We show that quantum effects generate a milder singularity at the position of the particle than the classical solution, but the singularity does not disappear. At large distances from the particle, the quantum correction decreases slowly, as $1/\rho^2$ (ρ being the distance to the particle in the shock wave plane). We argue that this large distance correction is a necessary consequence of quantum gravity.

04.50.+h, 04.30.Nk

Typeset using REVTeX

*Electronic Address: lousto@mail.physics.utah.edu

†Electronic Address: fmazzi@df.uba.ar

I. INTRODUCTION

The backreaction of quantum fields on the spacetime geometry can be described by the so called Semiclassical Einstein Equation (SEE) [1]

$$G_{\mu\nu} + \alpha H_{\mu\nu} + \beta I_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G \left(T_{\mu\nu}^{\text{class}} + \langle T_{\mu\nu} \rangle \right) \quad (1.1)$$

where one includes as a source the mean value of the energy momentum tensor of the quantum fields. The higher derivative terms

$$\begin{aligned} H_{\mu\nu} &= -2R_{;\mu\nu} + 2g_{\mu\nu}\square R - \frac{1}{2}g_{\mu\nu}R^2 + 2RR_{\mu\nu} \\ I_{\mu\nu} &= -2R_{\mu;\nu\alpha}^{\alpha} + \square R_{\mu\nu} + \frac{1}{2}g_{\mu\nu}\square R + 2R_{\mu}^{\alpha}R_{\alpha\nu} - \frac{1}{2}g_{\mu\nu}R_{\alpha\beta}R^{\alpha\beta} \end{aligned} \quad (1.2)$$

are needed to renormalize the theory. The dressed coupling constants $G, \Lambda, \alpha, \beta$ depend on an energy scale μ introduced by the regularization method. The dependence is given by the renormalization group equations. The mean value $\langle T_{\mu\nu} \rangle$ also depend on μ in such a way that the full equation is μ -independent. The numerical values of α and β should be determined by experiments or could be computed from a more fundamental theory. They should be of the order of M_{Plank}^2 for $\mu \sim M_{Plank}$.

Although a complete theory of quantum gravity from which the SEE could be derived is still lacking, there are several arguments that lead formally to this equation. On one hand, one can expand the metric around a classical solution and quantize both the linearized metric fluctuations and the matter fields on the classical background [1]. The one loop effective action for this field theory gives the leading correction (in powers of $\hbar$) to the classical Einstein equation. In this approximation, $\langle T_{\mu\nu} \rangle$ contains both graviton and matter contributions. As two loop corrections will induce additional terms proportional to $\hbar^2$, in this context it is reasonable to look only for solutions to the SEE which are perturbative in $\hbar$, and therefore close to the classical solution. On the other hand, the SEE can be derived using a $1/N$ approximation. Assume there are N decoupled quantum matter fields. Taking the limit $N \rightarrow \infty$ with GN fixed (and rescaling also the classical energy momentum tensor so that $GT_{\mu\nu}^{\text{class}}$ is N -independent for large N), one arrives [2] at an equation of the form (1.1). In this case, there is no graviton contribution to $\langle T_{\mu\nu} \rangle$, since it is suppressed by a factor $1/N$ with respect to the contribution of the N matter fields. Using this point of view, the SEE is valid to all orders in $\hbar$, and it makes sense to look for exact, self-consistent solutions. However, it should be kept in mind that some of the self-consistent solutions to the SEE may be non-physical. It is not clear that one can ignore graviton effects at Planck curvature, even for large N . Moreover, Eq. (1.1) contains time derivatives of order higher than two, and therefore ‘runaway’ solutions could arise. We refer the reader to Ref. [3] for discussions on this important issue.

Physical, exact solutions to the SEE are interesting because they can show important deviations from the classical behavior. The standard lore about smearing of classical singularities due to quantum effects should be proved in this context. The problem of the endpoint of Hawking radiation should, and in fact it is, being analyzed using the SEE, although progress has been important only in two dimensions [4]. In four dimensions, $\langle T_{\mu\nu} \rangle$ is a complicated, non-local function of the geometry, and the SEE has been solved in a few

cosmological cases [5]. In this paper we are going to find a plane fronted with parallel rays (pp) – gravitational shock wave self-consistent solution to the SEE. From the exact solution we will be able to discuss the problem of the smearing of the classical singularity. We will also compare the exact and perturbative solutions.

The paper is organized as follows. In Section II we introduce the pp–gravitational shock waves, and prove some useful geometric identities. In Section III we use a resummation of the Schwinger DeWitt expansion to write down the SEE for this particular metric. The SEE is solved in Section IV. We discuss our results in Section V and leave for the Appendix the case of the ultrarelativistic straight cosmic string.

II. THE GRAVITATIONAL SHOCK WAVE

It was stressed by 't Hooft [6] that at Planckian energies the relevant interaction among particles is the gravitational one, which takes a particular simple form. In fact the gravitational field generated by a highly energetic particle can be described by the metric of a gravitational shock wave

$$ds^2 = -du \, dv + f(x_\perp) \delta(u) \, du^2 + dx_\perp^2 , \quad (2.1)$$

where $u = t - z$ and $v = t + z$ are the usual null coordinates, and $dx_\perp^2$ is the two-dimensional flat metric. The wave propagates on a flat background along the trajectory of the particle, i.e. $u = 0$. The “profile” function $f(x_\perp)$ completely characterizes the spacetime.

Geodesics that cross the shock wave suffer a sudden shift in its v coordinate given by [7]

$$\Delta v = -f(\rho_i) , \quad (2.2)$$

where ρ_i is the distance to the inplunging geodesic from the source on the surface $u = 0$. This effect is represented in Fig. 1.

Geodesics also have a refraction effect given by [7]

$$\cot(\theta_{in}) + \cot(\theta_{re}) = \frac{1}{2} \partial_\rho f(\rho_i) , \quad (2.3)$$

where θ_{in} and θ_{re} are the incident and refracted angles respectively. This effect is depicted in Fig. 2.

The SEE (1.1) in principle involve terms containing powers of the curvature tensor, which in turn would be proportional to powers of the delta-function $\delta(u)$. These powers are meaningless in the context of classical distribution theory. In order to avoid this problem, we will consider a more general metric [13]

$$ds^2 = -du \, dv + H(x_\perp, u) \, du^2 + dx_\perp^2 . \quad (2.4)$$

We will show that for an arbitrary smooth function $H(x_\perp, u)$ the SEE becomes linear in the curvature and, at the end of the calculation, we will consider, the limit $H_n(x_\perp, u) \rightarrow f(x_\perp) \delta(u)$ ($\{H_n\}$ is a convergent sequence of well-behaved test functions).

The curvature scalars of the metric (2.4) formed out of the Ricci squared, Kretschmann, and Curvature scalar tensors, all identically vanish

$$R \equiv 0, \quad R_{\alpha\beta}R^{\alpha\beta} \equiv 0, \quad R_{\alpha\beta\gamma\delta}R^{\alpha\beta\gamma\delta} \equiv 0. \quad (2.5)$$

This does not mean, however, that the spacetime is free from any singularity. In fact, the non-vanishing components of the Riemann tensor are (apart from others obtained by symmetry properties)

$$R_{iuj u} = -\frac{1}{2}\partial_{ij}^2 H(x_\perp, u), \quad (2.6)$$

which may still diverge (as we will next see).

From Eq. (2.6) we see that the only non-vanishing component of the Ricci tensor is

$$R_{uu} = -\frac{1}{2}\nabla_\perp^2 H. \quad (2.7)$$

Note that the only non vanishing components of the inverse metric are g^{uv}, g^{vu}, g^{ii} and g^{vv} . Also that the Christoffel symbols with a subindex $v, \Gamma_{v\nu}^\mu$, vanish. This implies that the covariant derivative ∇_v coincides with the ordinary derivative when applied to a covariant index of a tensor, and vanishes when acting on a v -independent function.

From the above properties it is easy to show that *any scalar formed out of the Riemann tensor and its derivatives must vanish*. Indeed, a generic scalar can be built up as

$$g^{\mu\tau_1} g^{\nu\tau_2} \dots g^{\cdot\cdot} (\nabla_{\lambda_1} \nabla_{\lambda_2} \dots \nabla_{\lambda_n} R_{\mu\nu\rho\sigma}) (\nabla \dots \nabla R \dots) \dots (\nabla \dots \nabla R \dots). \quad (2.8)$$

The indices τ_i can be contracted with any of the covariant indices of the tensors in $(\nabla \dots \nabla R \dots)$. As the only non-vanishing components of the Riemann tensor are those given by Eq. (2.6), either μ or ν should equal u . Therefore, τ_1 or τ_2 should equal v . As $R_{v\dots} = 0$ and $\nabla_v R_{\dots} = 0$, the scalar (2.8) must vanish. Note that what we have shown is that any expression containing $R_{\mu\nu\rho\sigma}$ vanishes if both indices $\mu\nu$ (or, by symmetry, $\rho\sigma$) are contracted with any other index.

With similar arguments one can show that the most general tensor with two covariant indices must be a linear combination of $g_{\mu\nu}$ and $F(\nabla_\perp^2)R_{\mu\nu}$ [with F an arbitrary function].

The proof is as follows. Consider a generic tensor $(2, 0)$

$$g^{\mu\tau_1} g^{\rho\tau_2} \dots g^{\cdot\cdot} (\nabla_{\lambda_1} \nabla_{\lambda_2} \dots \nabla_{\lambda_n} R_{\mu\nu\rho\sigma}) (\nabla \dots \nabla R \dots) \dots (\nabla \dots \nabla R \dots) \quad (2.9)$$

The difference with respect to the scalar case is that now we can have up to two non-contracted indices in the Riemann tensor. If $R_{\mu\nu\rho\sigma}$ has three or four indices contracted, the previous arguments imply that the tensor must vanish. Therefore, $R_{\mu\nu\rho\sigma}$ must have exactly two non-contracted indices and, as we are considering a $(2, 0)$ tensor, it must be linear in $R_{\mu\nu\rho\sigma}$. For τ_1 and τ_2 there are two alternatives:

a) if τ_1 is contracted with τ_2 , the tensor becomes $\nabla_{\lambda_1} \nabla_{\lambda_2} \dots \nabla_{\lambda_n} R_{\nu\sigma}$. The indices λ_i must be contracted among themselves, and one finds something proportional to $(\nabla_\perp^2)^{\frac{n}{2}} R_{\nu\sigma}$.

b) if τ_1 and τ_2 are contracted with two derivatives λ_1 and λ_2 , the result is proportional to $\nabla_\perp^2 R_{\nu\sigma}$. The other derivatives must be contracted among themselves, and one finds again the same result.

The conclusion of this section is that any geometrical tensor built up from the metric, with two indices, and in particular, as we will soon see $\langle T_{\mu\nu} \rangle$ (when the state is the Minkowski's in-vacuum state), must be a linear combination of $g_{\mu\nu}$ and a function of $\nabla_\perp^2$ acting on $R_{\mu\nu}$.

III. THE SEMICLASSICAL EINSTEIN EQUATION

Let us write Eq. (1.1) for the particular case of the shock wave metric (2.1). Using the fact that $H_{\mu\nu} = 0$, $I_{\mu\nu} = \nabla_\perp^2 R_{\mu\nu}$, we find

$$R_{\mu\nu} + \beta \nabla_\perp^2 R_{\mu\nu} = 8\pi G \left(T_{\mu\nu}^{\text{class}} + F(\nabla_\perp^2) R_{\mu\nu} \right) \quad (3.1)$$

Here (as in Eq. (1.1)), we have split the stress–energy–momentum tensor into two pieces: a classical source plus the renormalized expectation value of the $T_{\mu\nu}$ due to N quantum matter fields. Since the state in which we take the expectation value is Minkowski’s vacuum, these fields do not contribute to the classical source and, since the fluctuations of the classical source are suppressed by a factor $1/N$ (as are the graviton contributions) neither of these two terms appear in (3.1) [For more details, see, for example, Refs. [8,9].]

In general, we expect the function F to depend on the mass of the quantum fields and on the scale μ introduced by the regularization method. For massive fields, this function can be expanded in powers of $(\nabla_\perp^2/m^2)$ (Schwinger–DeWitt expansion). For massless fields this expansion is inadequate, and F becomes a non-analytic function of $\nabla_\perp^2$. All this can be checked from the explicit calculations done by Vilkovisky and collaborators [10]. They computed the effective action up to quadratic terms in the curvature, and the energy momentum tensor up to linear terms in the curvature. But for the pp-shock wave we have shown this is an exact result! The energy momentum tensor *is* linear in the curvature. Therefore, the function F is given by [10,12]

$$F(\nabla_\perp^2) = \frac{m^2}{16\pi^2} \left(\xi - \frac{1}{6} \right) \ln \left[\frac{m^2}{\mu^2} \right] + \frac{\nabla_\perp^2}{384\pi^2} \int_0^1 dt \, t^4 \ln \left[\frac{m^2 - \frac{1}{4}(1-t^2)\nabla_\perp^2}{\mu^2} \right] . \quad (3.2)$$

As anticipated, if $m^2 \neq 0$, it is possible to expand F in powers of $(\nabla_\perp^2/m^2)$

$$F(\nabla_\perp^2) = \frac{m^2}{16\pi^2} \left(\xi - \frac{1}{6} \right) \ln \left[\frac{m^2}{\mu^2} \right] + \frac{\nabla_\perp^2}{384\pi^2} \int_0^1 dt \, t^4 \left[\ln \left[\frac{m^2}{\mu^2} \right] - \frac{1}{4}(1-t^2) \frac{\nabla_\perp^2}{m^2} + \mathcal{O} \left(\left(\frac{\nabla_\perp^2}{m^2} \right)^2 \right) \right] , \quad (3.3)$$

the Schwinger DeWitt expansion is recovered.

On the other hand, if $m^2 = 0$, the form factor reads

$$F(\nabla_\perp^2) = \frac{\nabla_\perp^2}{1920\pi^2} \ln \left[-\frac{\nabla_\perp^2}{\mu^2} \right] + \text{const} \, \nabla_\perp^2 \quad (3.4)$$

and it is a non-analytic function of the Laplacian.

It is interesting to note that in the massless case the form of the function F can be derived using general arguments [11]. Indeed, from dimensional analysis we know that F must be of the form $F(\mu^2, \nabla_\perp^2) = \nabla_\perp^2 \tilde{F}(\nabla_\perp^2/\mu^2)$. Moreover, G is independent of μ and β depends on μ according to the renormalization group equation [12]

$$\frac{\mu}{8\pi G} \frac{d\beta}{d\mu} = -\frac{1}{960\pi^2} . \quad (3.5)$$

As the whole semiclassical equation must be independent of μ , we must have $\mu d\tilde{F}/d\mu = -1/(960\pi^2)$. This equation fixes the function F up to a term that can be absorbed into a redefinition of β , and the result coincides with Eq. (3.4).

Now we are ready to write the differential equation for the profile function. Assuming a point-like classical source ¹

$$T_{uu}^{\text{class}} = p\delta(x_\perp)\delta(u) \quad (3.6)$$

we obtain

$$-\frac{1}{2}\left[\nabla_\perp^2 + \left(\beta(\mu) - \mathcal{A}\ln\left(-\frac{\nabla_\perp^2}{\mu^2}\right)\right)\nabla_\perp^4\right]f(\rho) = 8\pi Gp\delta(x_\perp); \quad \mathcal{A} = \frac{G}{240\pi} \quad (3.7)$$

Note that the μ -independence of the above equation is explicit: Eq.(3.5) implies that the parenthesis multiplying $\nabla_\perp^4$ in Eq. (3.7) is independent of μ . Eq.(3.7) has been solved in Ref. [13] for the case $\mathcal{A} = 0$ (i.e. for the fourth order theory).

IV. SOLUTION TO THE SEMICLASSICAL EQUATION

Plane gravitational shock waves (2.1) have the nice property of linearizing field equations. This feature allowed us to write Eq. (3.7) in an explicit form. Again, this linearization allow us to seek for the solution to the field equation by using Fourier transform methods. The result is

$$f(x_\perp) = f_{GR} + 16\pi Gp \int \frac{d\vec{k}}{4\pi^2} e^{i\vec{k}\cdot\vec{x}_\perp} \frac{[\beta - \mathcal{A}\ln(\frac{k^2}{\mu^2})]}{[1 - \beta k^2 + \mathcal{A}k^2 \ln(\frac{k^2}{\mu^2})]}, \quad (4.1)$$

where $f_{GR} = -8Gp\ln(\rho/\rho_0)$ is the general relativistic result [14], with ρ_0 an arbitrary constant. The second term in the above equation contains the corrections to the classical result.

To perform this integral we first note that since the problem has axial symmetry it is convenient to pass from Cartesian to polar coordinates in the perpendicular space, i.e. from $x_\perp$ to (ρ, φ) . Doing so, the angular integral in Eq. (4.1) is a representation of the Bessel function [15] J_0

$$f(\rho) = f_{GR} + 8Gp \int_0^\infty dk \, k J_0(k\rho) \frac{[\beta - \mathcal{A}\ln(\frac{k^2}{\mu^2})]}{[1 - \beta k^2 + \mathcal{A}k^2 \ln(\frac{k^2}{\mu^2})]}. \quad (4.2)$$

Note that in the fourth order theory ($\mathcal{A} = 0$, $\beta \neq 0$), the integrand above has no real poles when the so called non-tachyon constraint, $\beta < 0$, is satisfied. When $\mathcal{A} \neq 0$, the condition for absence of real poles is $\mathcal{A}\mu^2 \exp(\beta/\mathcal{A}) < e$.

¹This is an approximation, but if one wants to deal with an extended source, our results below give the Green's function for that case.

In this integral (now on the magnitude of $\vec{k}$), if $\mathcal{A} \neq 0$ we can absorb the term proportional to β into a redefinition of the scale μ

$$f(\rho) = f_{GR} - 8Gp \int_0^\infty dk \frac{\mathcal{A} \ln(k^2 \mu^{-2} e^{-\beta/\mathcal{A}})}{1 + \mathcal{A} k^2 \ln(k^2 \mu^{-2} e^{-\beta/\mathcal{A}})} k J_0(k\rho) . \quad (4.3)$$

Let us now discuss some properties of the self-consistent profile function. We first note that it is finite at the origin $\rho = 0$. Indeed, we can write

$$\begin{aligned} f(\rho) &= -8pG \left[\ln(\rho/\rho_0) + K_0(\rho/\sqrt{\mathcal{A}}) - \int_0^\infty d\bar{k} J_0(\bar{k}\rho^*) \left(\frac{\bar{k} \ln\left(\frac{\bar{k}^2}{\mathcal{A}^*}\right)}{1 + \bar{k}^2 \ln\left(\frac{\bar{k}^2}{\mathcal{A}^*}\right)} - \frac{\bar{k}}{1 + \bar{k}^2} \right) \right] \\ &\doteq f_{GR} + f_Q + \Delta f . \end{aligned} \quad (4.4)$$

where we have taken dimensionless variables $\bar{k} \doteq \sqrt{\mathcal{A}}k$, $\rho^* \doteq \rho/\sqrt{\mathcal{A}}$ and $\mathcal{A}^* \doteq \mathcal{A}\mu^2 \exp(\beta/\mathcal{A})$ for $\mathcal{A} \neq 0$. Also, we identified with the subindices “GR”, the general relativistic result, with “Q” the corrections induced by terms quadratic in the curvature (proportional to K_0), and with Δf the higher order corrections.

Using the asymptotic behavior of the modified Bessel function K_0 and noting that the last integral in Eq. (4.4) is finite in the limit $\rho \rightarrow 0$ one can easily check that $f(\rho)$ is finite in this limit. This confirms the results of Ref. [13] where the first two addends of Eq. (4.4) have been obtained.

The next question is about the singularity at the origin. Although all curvature scalars vanish for the pp-wave, the curvature tensor itself may diverge. In fact,

$$R_{\rho u \rho}^v = \delta(u) \partial_\rho^2 f(\rho) , \quad R_{\phi u u}^\phi = -\frac{\delta(u)}{2\rho} \partial_\rho f(\rho) , \quad R_{\rho u u}^\rho = \frac{\delta(u)}{2} \partial_\rho^2 f(\rho) , \quad (4.5)$$

that for the classical solution diverge as $\partial_\rho^2 f_{GR} \sim \rho^{-1} \partial_\rho f_{GR} \sim \rho^{-2}$. When quadratic terms in the curvature are included in the classical action, the divergence is milder, the components of the curvature tensor diverge logarithmically [13] $\partial_\rho^2 (f_{GR} + f_Q) \sim \rho^{-1} \partial_\rho (f_{GR} + f_Q) \sim \ln \rho$. We will now show that the divergence is still milder for the self-consistent solution. The second derivative of the profile function can be written as

$$\partial_\rho^2 f = 8pG \int_0^\infty d\bar{k} \frac{\bar{k} J_0''(\bar{k}\rho^*)}{1 + \bar{k}^2 \ln \frac{\bar{k}^2}{\mathcal{A}^*}} \quad (4.6)$$

where the primes denote derivatives with respect to the argument. Note that since $J_0''(0) = -1/2$, this integral diverges as the $\ln[\ln(\cdot)]$ of the upper limit.

To see the dependence with ρ we split this integral as

$$\partial_\rho^2 f \simeq 8pG \left[\int_0^{\bar{k}_0} d\bar{k} \frac{\bar{k} J_0''(\bar{k}\rho^*)}{1 + \bar{k}^2 \ln \frac{\bar{k}^2}{\mathcal{A}^*}} + \int_{\bar{k}_0}^\infty d\bar{k} \frac{J_0''(\bar{k}\rho^*)}{\bar{k} \ln \frac{\bar{k}^2}{\mathcal{A}^*}} \right] \quad (4.7)$$

where $\bar{k}_0$ is such that $\bar{k}_0^2 \ln \frac{\bar{k}_0^2}{\mathcal{A}^*} \gg 1$. The first term in Eq. (4.7) is finite in the limit $\rho \rightarrow 0$. The second term gives, after introducing the new variable $y = \bar{k}\rho^*$ and performing an integration by parts

$$\begin{aligned}
\int_{\bar{k}_0\rho}^{\infty} \frac{dy}{y} \frac{J_0''(y)}{\ln \frac{y^2}{\mathcal{A}^* \rho^{*2}}} &= \frac{1}{2} J_0''(y) \ln \left[\ln \left(\frac{y^2}{\mathcal{A}^* \rho^{*2}} \right) \right] \Big|_{\bar{k}_0\rho^*}^{\infty} - \frac{1}{2} \int_{\bar{k}_0\rho^*}^{\infty} dy J_0'''(y) \ln \left[\ln \left(\frac{y^2}{\mathcal{A}^* \rho^{*2}} \right) \right] \\
&\simeq \frac{1}{2} J_0''(\bar{k}_0\rho^*) \left\{ \ln \left[\ln \left(\frac{1}{\mathcal{A}^* \rho^{*2}} \right) \right] - \ln \left[\ln(\bar{k}_0/\mathcal{A}^*) \right] \right\} .
\end{aligned} \tag{4.8}$$

Using that $J''(0) = -1/2$, we see that $\partial_\rho^2 f \simeq 2pG \ln[-\ln(\rho^*)]$ as $\rho \rightarrow 0$. A similar result can be shown for $\rho^{-1} \partial_\rho f$. The conclusion is that the curvature diverges like $\ln[-\ln \rho^*]$ near the position of the particle.

What happens at large distances? In this case we expect the self-consistent solution to approach the solution perturbative in $\hbar$ (perturbative in $\mathcal{A}$, in our notation). From Eq. (4.4) we see that, to lowest order in $\mathcal{A}$ (for large ρ),

$$\Delta f(\rho) \simeq -8pG \int_0^\infty d\bar{k} \bar{k} J_0(\bar{k}\rho^*) \ln \left(\frac{\bar{k}^2}{\mathcal{A}^*} \right) . \tag{4.9}$$

This integral is ill defined and must be treated as a distribution. It is essentially the Fourier transform of $\ln(\bar{k}^2/\mathcal{A}^*)$ in two dimensions. To compute it, we introduce an integral representation for the logarithm

$$\Delta f(\rho) \simeq 8Gp \int_0^\infty dz \int_0^\infty d\bar{k} \bar{k} J_0(\bar{k}\rho^*) \left[\frac{1}{z + \bar{k}^2} - \frac{1}{z + \mathcal{A}^*} \right] . \tag{4.10}$$

The $\mathcal{A}^*$ -dependent term gives a contribution proportional to $\delta(x_\perp)$ which is irrelevant at large distances. Omitting this term we obtain

$$\Delta f(\rho) \simeq 8Gp \int_0^\infty dz K_0(\sqrt{z}\rho^*) = \frac{16Gp\mathcal{A}}{\rho^2} . \tag{4.11}$$

Therefore, we expect that, at large distances, $f(\rho) = f_{GR} + 16Gp\mathcal{A}/\rho^2$. It is important to stress that the perturbative and self-consistent solutions are completely different unless $\rho \rightarrow \infty$.

We have numerically computed the self-consistent solution starting from Eq. (4.4), and confirmed the properties of the profile function both at the $\rho \rightarrow 0$ and $\rho \rightarrow \infty$ limits. The results for Δf and $\partial_\rho^2 f$ are shown in Figs. (3) and (4). In those figures we only show curves for $\mathcal{A}^* < e$, since Δf diverges for $\mathcal{A}^* = e$. This is due to the fact that, as we already pointed out, the integrand in Eq. (4.1) develops a double real pole as $\mathcal{A}^*$ approaches e , that becomes two single real poles for $\mathcal{A}^* > e$. These two real poles, k_p , change the asymptotic behavior of Δf for large ρ^* , making it to decrease slowly (like a linear combination of $J_0(k_p\rho)$, $p = 1, 2$). However, from the definition of $\mathcal{A}^*$ right after Eq. (4.4), we and the renormalization group equation (3.5), we see that $\mathcal{A}^*$ is actually independent of μ . Then, from the renormalized value of $\mathcal{A}$, given in Eq. (3.7), and the no tachyon constraint that implies $\beta < 0$, we can infer that $\mathcal{A}^* \ll 1$.

V. DISCUSSION

We have found a self-consistent pp-shock wave solution to the SEE in the $1/N$ approximation.

We have shown that quantum effects make milder the singularity of the classical solution at the origin. In the classical theory, the curvature diverges like $1/\rho^2$. In the quadratic theory it diverges like $\ln(\rho^2)$. When the backreaction effects of massless quantum scalar fields are taken into account, it diverges like $\ln[-\ln(\rho^2)]$. Therefore, the self-consistent solution can be trusted up to a distance very close to the position of the particle, where the curvature becomes of the order of Planck curvature.

At large distances, the situation is similar to the case of the corrections to the Newtonian potential: while classical terms quadratic in the curvature produce an exponentially decreasing correction [16]

$$V = -\frac{GM}{r}(1 - \exp(-2\sigma r)) , \quad \sigma^2 = 3\alpha + \beta , \quad (5.1)$$

backreaction effects due to massless quantum fields induce a power law correction of the form [12,17]

$$V = -\frac{GM}{r} + \frac{aG^2M}{r^3} . \quad (5.2)$$

This is the dominant correction at large distances. The numerical constant a depends on the matter content of the theory and includes a contribution from gravitons.

We have a similar picture for the pp-gravitational shock wave. The profile function in the quadratic theory is given by [13]

$$f_{GR} + f_Q = f_{GR}(\rho) - 8GpK_0(\rho/\sqrt{-\beta}) , \quad (5.3)$$

and the correction to the GR result is exponentially small at large distances. However, we have shown that quantum massless fields induce a correction of the form

$$f(\rho) \simeq f_{GR}(\rho) + \frac{16Gp\mathcal{A}}{\rho^2} ; \quad \rho \rightarrow \infty , \quad (5.4)$$

which again is the leading correction at large distances.

If one treats General Relativity as an effective field theory of an unknown quantum theory of gravity, it can be shown that the leading quantum corrections in Eq. (5.2) will not depend on the details of the more fundamental theory: they are a consequence of quantum gravity, whatever this theory may be [17]. The same argument applies to the profile function of the gravitational shock wave: we expect quantum gravity effects to produce large distance corrections proportional to $\mathcal{C}/\rho^2$. The coefficient $\mathcal{C}$ should be determined by a more careful calculation that should take into account all massless fields in nature including graviton contributions to the renormalized energy-momentum tensor. However, this contribution should be qualitatively similar to the one computed for massless scalar fields in this paper.

Since we are considering exact solutions to the semiclassical Einstein equations (1.1), it is worth discussing here the question of the runaway solutions that such higher order equations may have. In Ref. [18] it was proposed that a way of getting rid of such unwanted runaway solutions is to only consider first order perturbative solutions in the corrections to general relativity (here proportional to $\mathcal{A}$). This is clearly too restrictive in the framework presented in our paper. In the $1/N$ approach ($N \rightarrow \infty$) the semiclassical Einstein equations

hold exactly. In fact, since the scalar fields are taken to be free (i.e. non-self-interacting) only the leading order terms in these scalar fields arise. Consequently, one is not only allowed to go further than the first order perturbative regime, but one should also take into account all the “well-behaved” exact solutions to the semiclassical fields equations. The well-behaved solutions we found have been selected from all possible ones by allowing them to be represented by a Fourier transform.

ACKNOWLEDGMENTS

C.O.L. was partially supported by NSF Grant No. PHY-95-07719 and by research funds of the University of Utah. F.D.M. was supported by Universidad de Buenos Aires, CONICET and Fundación Antorchas. The authors thank M. Campanelli for useful discussions.

APPENDIX A: ULTRARELATIVISTIC COSMIC STRING

It is also of interest to consider the case of an ultrarelativistic particle in 2+1 dimensions, since its results are also valid for a straight cosmic string boosted in a direction perpendicular to it [19,13].

In this case the field equation simply takes the form of Eq. (3.7) with $\nabla_{\perp}^2$ replaced by ∂_y^2 , where y is the Cartesian coordinate perpendicular to both the direction of motion and the axis along the string.

Consequently, the Fourier decomposition of the profile function leads to

$$f(|y|) = f_{GR}(|y|) + 16\pi G\tilde{p} \int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{ik|y|} \frac{\left[\beta - \mathcal{A} \ln\left(\frac{k^2}{\mu^2}\right)\right]}{\left[1 - \beta k^2 + \mathcal{A} k^2 \ln\left(\frac{k^2}{\mu^2}\right)\right]}, \quad (\text{A1})$$

where now $\tilde{p}$ represents the momentum per unit length of the cosmic string and $f_{GR} = -8\pi\tilde{p}|y|$ is the general relativistic result found in Ref. [19].

By splitting the integral in Eq.(A1) we obtain

$$f(|y|) = f_{GR}(|y|) + 8G\tilde{p} \left\{ -2\sqrt{\mathcal{A}} \int_0^{\infty} d\bar{k} \frac{\cos(\bar{k}|y|^*)}{1 + \bar{k}^2} + 2\sqrt{\mathcal{A}} \int_0^{\infty} d\bar{k} \cos(\bar{k}|y|^*) \left[\frac{1 - \ln(\bar{k}^2/\mathcal{A}^*)}{(1 + \bar{k}^2)(1 + \bar{k}^2 \ln(\bar{k}^2/\mathcal{A}^*))} \right] \right\}, \quad (\text{A2})$$

where the dimensionless variables have the same meaning as given after Eq. (4.4). The first integral can be found in tables [15] and its result is $\pi \exp\{-|y|/\sqrt{\mathcal{A}}\}/2$. Thus giving the quadratic theory correction as found in Ref. [13]. The second integral has to be performed numerically as we did before for the particle in four dimensions.

It is easy to check that

$$f''(|y|) = \frac{-16G\tilde{p}}{\sqrt{\mathcal{A}}} \int_0^{\infty} d\bar{k} \frac{\cos(\bar{k}|y|^*)}{1 + \bar{k}^2 \ln(\bar{k}^2/\mathcal{A}^*)}, \quad (\text{A3})$$

is finite at $|y| = 0$. The Riemann tensor is then also finite (apart from the $\delta(u)$ behavior). Note that this is already true in the quadratic theory; while in general relativity, the divergence was as $\delta(|y|)$.

The behavior for large $|y|$ can be computed on the same lines as before, and we obtain

$$f(|y|) \simeq -8\pi G\tilde{p}|y| + \frac{16\pi G\tilde{p}\mathcal{A}}{|y|} ; \quad |y| \rightarrow \infty . \quad (\text{A4})$$

Once more, we see that quantum effects due to massless fields are the leading corrections at large distances.

REFERENCES

- [1] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press, Cambridge (1982).
- [2] J. B. Hartle and G. T. Horowitz, Phys. Rev. D **24**, 257 (1981).
- [3] E. E. Flanagan and R. M. Wald, Phys. Rev. D **54**, 6233 (1996), and references therein.
- [4] A. Strominger, "Les Houches Lectures on Black Holes", hep-th 9501171, and references therein.
- [5] M. V. Fischetti, J. B. Hartle and B. L. Hu, Phys. Rev. D **20**, 1757 (1979); J.B. Hartle and B. L. Hu, ibid 1772 (1979); A. A. Starobinsky, Phys. Lett. **B** 91, 99 (1980); P. Anderson, Phys. Rev. D **29**, 615 (1984).
- [6] G. 't Hooft, Phys. Lett. B **198**, 61 (1987).
- [7] T. Dray and G. 't Hooft, Nucl. Phys. B **253**, 173 (1985).
- [8] M. Campanelli, Ph. D. Thesis, University of Bern, June 1996 (unpublished).
- [9] M. Campanelli and C. O. Lousto, preprint gr-qc/9608027.
- [10] A. O. Barvinsky and G. A. Vilkovisky, Nucl. Phys. B **282**, 163 (1987); ibid B **333**, 471 (1990); G. A. Vilkovisky in *Quantum Theory of Gravity*, ed. by S.M. Christensen (Hilger Bristol, 1984).
- [11] F.C. Lombardo and F.D. Mazzitelli, preprint gr-qc/9609073.
- [12] D. A. R. Dalvit and F. D. Mazzitelli, Phys. Rev. D **50**, 1001 (1994).
- [13] M. Campanelli and C. O. Lousto, Phys. Rev. D **54**, 3854 (1996).
- [14] P. C. Aichelburg and R. U. Sexl, Gen. Rel. Grav. **2**, 303 (1971).
- [15] I. S. Gradshteyn and I. M. Ryzhik, *Tables of integrals, Series and Products*, Academic Press, New York (1980).
- [16] K.S. Stelle, Gen. Rel. and Grav. **9**, 353 (1978).
- [17] J.F. Donoghue, Phys. Rev. D **50**, 3874 (1994).
- [18] L. Parker and J. Z. Simon, Phys. Rev. D, **47**, 1339 (1993).
- [19] C. O. Lousto and N. Sánchez, Nucl. Phys. B **355**, 231 (1991).


FIGURES

FIG. 1. The dashed line represents the path of a null geodesics in the (u, v) plane. Since for $u \neq 0$ the spacetime is flat, geodesics are straight lines. At $u = 0$ they have a discontinuity jump described by Eq. (2.2). The point-like source p is located at the origin and the plane $u = 0$ contains the gravitational shock wave.

FIG. 2. The ‘spatial refraction’ of a null geodesics as described by Eq. (2.3). Here θ_{in} and θ_{re} are the incident and refracted angles respectively.

FIG. 3. The higher order corrections to the profile function of the shock wave as defined in Eq. (4.4) in units of $8pG$. Here $\rho^* = \rho/\sqrt{\mathcal{A}} \cong 27.5l_{Pl}^{-1}\rho$ and $\mathcal{A}^* \doteq \mathcal{A}\mu^2 \exp(\beta/\mathcal{A})$ are convenient dimensionless variables. The correction reaches a finite value at $\rho = 0$ and vanishes as $1/\rho^{*2}$ for large ρ^* .

FIG. 4. $d^2f/d\rho^2$ as defined in Eq. (4.6) over $8pG$. This is proportional to the non-vanishing Riemann tensor components and show that they diverge as $\ln[\ln(\rho^*)]$ for $\rho^* \rightarrow 0$.






Figure 3




Figure 4

